



PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:)

KUNDIG and McCORMACK)

Serial No.: 09/804,464)

Filed: March 13, 2001)

Group Art Unit: 1644

Examiner: Phyoung N. HUYNH

Docket No: 005184.00002

For: **MODULATION OF ALLERGIC RESPONSE**

DECLARATION OF DR. THOMAS KÜNDIG UNDER 37 C.F.R. § 1.132

Commissioner of Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

I, Thomas Kündig, declare as follows:

1. I am named as an inventor of the subject matter claimed in the application referenced above.
2. I hold an M.D. degree from the University of Zürich and have worked in the fields of immunology and allergy for 14 years. My curriculum vitae is attached as Exhibit 1.
3. We conducted a clinical study that included a total of 153 human hay fever sufferers. Each of these patients was allergic to grass pollen, as assessed by skin prick testing to grass pollen extract. Patients were randomized into two study groups. Eighty-six patients in the conventional treatment group received conventional grass pollen immunotherapy. These patients were subcutaneously injected with 16 incremental doses of grass pollen extract over 20 weeks, followed by twice-weekly subcutaneous injections of a maintenance dose corresponding to 100

µg alum-absorbed grass pollen extract (100,000 so called "SQ-E units"). Sixty-seven patients in the intralymphatic group received three injections of 1 µg of the same alum-absorbed grass pollen extract into a superficial subcutaneous lymph node of the groin spaced four weeks apart. The efficacy of both the conventional and the intralymphatic treatments were assessed four months after the initial treatment. The protocol is shown in Exhibit 2.

4. To assess efficacy, patients in both treatment groups were evaluated for their sensitivity to the allergen by nasal provocation with an extract of the same pollen extract. Fifty µl of pollen extract were administered under rhinoscopic control onto the mucosa of the lower concha. A first dose of 100 SQ-E (approximately 0.1 µg of grass pollen extract) was followed after 30 minutes by 1,000 SQ-E (approximately 1 µg), then by 10,000 SQ-E (approximately 10 µg) and finally by 100,000 SQ-E (approximately 100 µg) after 60 and 90 minutes, respectively. The following symptoms and scores were recorded: ocular or nasal itching (0 = none; 1 = mild, *i.e.*, slight sensation; 2 = moderate, *i.e.*, definite sensation; and 3 = severe, *i.e.*, need to rub nose or eyes); runny nose (0= none; 1 = mild, *i.e.*, slight sensation; 2= moderate, *i.e.*, definite sensation; and 3 = severe, *i.e.*, need to blow or wipe nose); nasal congestion (0= none; 1= mild, *i.e.*, slight sensation; 2 = moderate, *i.e.*, definite sensation; and 3= severe, difficult to breathe through one or both nares); and sneezing (0= none; 1= mild, *i.e.*, 1-2 sneezes; 2 = moderate, *i.e.*, 3-4 sneezes; and 3= severe, *i.e.*, 5 or more sneezes). The results are shown in Exhibit 3.

5. The upper left panel of Exhibit 3 depicts the symptom scores observed after nasal provocation with 100 SQ-E units (approx. 0.1 µg of pollen extract). At this low pollen concentration the intranodal group shows fewer symptoms after treatment, but due to the generally low symptom scores at this low pollen concentration, the latter improvement of the intranodal group shows no statistical significance ($p= 0.45$). The upper right panel of Exhibit 3

shows the symptom scores after nasal provocation with 1,000 SQ-E units (approximately 1 µg of pollen extract). Again the intranodal group of patients shows a lower symptom score after treatment, but again the overall symptom scores are low, so that the improvement is not statistically significant ($p=0.07$).

6. The lower left panel of Exhibit 3 shows the symptom scores after nasal provocation with 10,000 SQ-E units (approximately 10 µg of pollen extract). Because of the higher pollen concentration, the overall symptom scores are higher and the improvement of the intranodal group becomes statistically significant (average score before treatment 4.47, compared with 1.9 after treatment). The lower right panel of Exhibit 3 shows the symptom scores after nasal provocation with 100,000 SQ-E grass pollen extract (approximately 100 µg pollen extract). Before treatment the intranodal group showed an average symptom score of 9.63; after treatment, the symptom score is reduced to 5.05. This change is statistically highly significant ($p=0.002$).

7. Thus, intralymphatic administration of allergen administration is more efficient than subcutaneous administration of the same allergen. No more than three intranodal injections of an allergen are sufficient to desensitize human patients to the allergen, as assessed by sensitivity to the allergen. This permits a reduced number of injections and therefore substantially shortens the duration of treatment.

8. Patients that were desensitized by only 3 intranodal injections were assessed by nasal provocations tests 4 months after the beginning of treatment. As shown in Exhibit 4, this patient group tolerated approximately 10 times higher pollen concentrations after treatment. In contrast, subcutaneously desensitized patients showed no significant improvement of their nasal provocation thresholds. In order to achieve a 10-fold increase in the nasal provocation threshold

conventional subcutaneous immunotherapy usually has to be continued for 3 years, as demonstrated by Wihl *et al.* for tree pollen (*Allergy* 43, 363-69, 1988; Exhibit 6). Similarly, Pegelow *et al.* (*Allergy* 39, 275-90, 1984; Exhibit 7) showed that after 3 years of immunotherapy with grass pollen the average increase in the nasal provocation threshold is not more than a factor of 10.

9. It is unexpected that treatment with merely three injections of 1 µg more efficiently reduces the symptom score than the extensive conventional subcutaneous regimen with 20 incremental injections up to 100 µg of the same pollen extract. Based on these results, I would expect that the efficiency of desensitization with other allergens such as venoms, animal dander, and dust mite would be similarly enhanced by intranodal administration.

10. Pollen-specific IgG also was measured before treatment and four months after the beginning of treatment. No significant changes were observed. Amongst all IgG subclasses, subclass IgG4, if at all, is the one subclass where increases have been reported by other groups. The graph shown in Exhibit 5 depicts specific IgG4 against the major allergenic grass in Europe (*i.e.*, *Phleum pratensae*) in patients treated by intralymphatic allergen injections. IgG4 titers were measured by the Pharmacia CAP™ method, which permits a quantitative assessment. There was no significant titer change in *Phleum pratensae*-specific IgG4 before and after treatment. Thus, the mechanism of intralymphatic desensitization does not depend on induction of allergen-specific IgG4 antibodies.

11. A total of 20 systemic allergic adverse events were observed during a one hour post-injection wait for the conventional group. Two of these events (asthma attacks) were classified as severe (grade 3 according to H.L. Müller, *New Engl. J. Med.* 261, 374-77, 1959; *J. Asthma Res.* 3, 331-33, 1966). Eighteen of these events were classified as mild or moderate

(grades 1 and 2 according to H.L. Müller); most of these patients suffered from flush, urticaria, or angioedema. In contrast, only six systemic allergic events were observed during a two hour post-injection wait for the intralymphatic group. These patients suffered from flush, urticaria or angioedema (grades 1 and 2 according to H.L. Müller). This demonstrates that the frequency and severity of adverse events are reduced with intralymphatic treatment compared to conventional treatment.

12. At the one year follow up visit, 40 of the original 86 patients in the conventional treatment group had dropped out of the study. In contrast, only three of the original 67 patients in the intralymphatic treatment group had dropped out. Thus, patients receiving intralymphatic treatment show better compliance and are more likely to finish their treatment.

13. I declare that all statements made herein of my own knowledge are true and that I believe all statements made on information and belief are true and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

14.10.03

Date

T. Kündig

Thomas Kündig, M.D.

Curriculum vitae of Thomas Kündig

Name: Kündig
First Name, Middle Name: Thomas, Martin
Titels: PD Dr.med.
Birth data: 22.1.1963
Nationality: Schwitzerland and USA
Working place: Dermatologische Klinik
Gloriastr. 31
8091 Zürich
Tel: 0041-1-255-3973
FAX: 0041-1-255-4418

Achievements:

- 1981 Graduation
Mathematical-Scientific type (Typus C),
Kantonsschule Zürcher Oberland
- 1987 MD degree
University of Zürich
- 1988 Medical Thesis
in respiratory physiology, supervisor Prof. E.A. Koller
University of Zürich
- 1988 Educational Commission for Foreign Medical Graduates
Basic Sciences, Clinical Sciences and English
Bern
- 1991 Post-graduate Course in Experimental Medicine
especially Immunology, supervisor Prof. R.M. Zinkernagel
Universität Zürich
- 1998 Venia legendi in Experimental Immunology and Experimental Dermatology

Working Places:

Kreisspital Wetzikon Dr. R. Stahel	Resident	1988-89	Surgery
Inst. for Experimental Immunology Universität Zürich Prof. R.M. Zinkernagel	Post-doc	1989-92	Immunology
Dept. of Medical Biophysics University of Toronto Prof. T.W.Mak	Post-doc	1992-95	Immunology
Dermatologische Klinik Universität Zürich Prof. G. Burg	Oberassistent Oberarzt	1995-99 1999-	Dermatology

Awards and Honors:

Award for best Marks at Graduation Exams 1987
Fellowship for Post-graduate Course in Experimental Medicine 1989-91.
Fellowship as an advanced researcher by Swiss National Science Foundation 1992-94.
Venia legendi in Experimental Immunology and Experimental Dermatology 1998
Georg-Friedrich Götz Award by the University of Zürich 1999 for outstanding medical research
on the development of a new method to enhance the immunogenicity of vaccines
Affiliated Researcher of the Ludwig Institute for Cancer Research

Publications (Original):

Boutellier, U. , **T. M. Kündig**, and E. A. Koller
Delay variations in automatic respiratory analysis systems.
Pflügers Archiv (1986) 234:6.

Boutellier, U. , **T. M. Kündig**, U. Gomez, P. Pietsch, and E. A. Koller. 1987. Respiratory phase detection and delay determination for breath-by-breath analysis.
J. Appl. Physiol. (1987) 62:837.

Binder, D. and **T. M. Kündig**
Antiviral protection by CD8⁺ versus CD4⁺ T cells: CD8⁺ T cells correlating with cytotoxic activity in vitro are more efficient in antivaccinia virus protection than CD4-dependent interleukins.
J. Immunol. (1991) 146:4301.

Rahemtulla, A. , W. P. Fung-Leung, M. W. Schilham, **T. M. Kündig**, S. R. Sambhara, A. Narendran, A. Arabian, A. Wakeham, C. J. Paige, R. M. Zinkernagel, R. G. Miller, and T. W. Mak
Normal development and function of CD8⁺ cells but markedly decreased helper cell activity in mice lacking CD4.
Nature (1991) 353:180.

Fung-Leung, W. P., **T. M. Kündig**, R. M. Zinkernagel, and T. W. Mak
Immune response against lymphocytic choriomeningitis virus infection in mice without CD8 expression.
J. Exp. Med. (1991) 174:1425.

Fung-Leung, W. P., M. W. Schilham, A. Rahemtulla, **T. M. Kündig**, M. Vollenweider, J. Potter, W. van Ewijk, and T. W. Mak
CD8 is need for development of cytotoxic T cells but not helper T cells.
Cell (1991) 65:443.

Althage, A. , B. Odermatt, D. Moskophidis, **T. Kündig**, H. Hengartner, and R. M. Zinkernagel
Immunosuppression by lymphocytic choriomeningitis virus infection: competent effector T and B cells but impaired antigen presentation.
Eur. J. Immunol. (1992) 22:1803.

Kündig, T. , A. Althage, H. Hengartner, and R. M. Zinkernagel
A skin test to assess virus specific cytotoxic T cell activity.
Proc. Natl. Acad. Sci. USA (1992) 89:7757.

Kündig, T. M., H. Hengartner, and R. M. Zinkernagel
T cell dependent IFN-gamma exerts an antiviral effect in the central nervous system but not in peripheral solid organs.

J. Immunol. (1993) 150:2316.

Oehen, S. , H. P. Waldner, **T. M. Kündig**, H. Hengartner, and R. M. Zinkernagel
Cytotoxic T cell memory to lymphocytic choriomeningitis virus is governed by elevated CTL-P frequencies and persisting antigen.
J. Exp. Med. (1993) 176:1273.

Castelmur, I. , C. DiPaolo, M. F. Bachmann, H. Hengartner, R. M. Zinkernagel, **T. M. Kündig**
Comparison of the sensitivity of in vivo and in vitro assays for detection of antiviral cytotoxic T cell activity.
Cell. Immunol. (1993) 151:460.

Kündig, T. M., H. Schorle, M. F. Bachmann, H. Hengartner, R. M. Zinkernagel, and I. Horak.
Immune responses in IL-2 deficient mice.
Science (1993) 262:1059.

Shahinian, A. , K. Pfeffer, K. P. Lee, **T. M. Kündig**, K. Kishihara, A. Wakeham, K. Kawai, P. S. Ohashi, C. B. Thompson, and T. W. Mak
Differential costimulatory requirements in CD28-deficient mice.
Science (1993) 261:609.

Schilham, M. W., W. P. Fung-Leung, A. Rahemtulla, **T. M. Kündig**, L. Zhang, J. Potter, R. G. Miller, H. Hengartner, and T. W. Mak
Alloreactive cytotoxic T cells can develop and function in mice lacking both CD4 and CD8.
Eur. J. Immunol. (1993) 23:1299.

Bachmann, M. F., **T. M. Kündig**, C. P. Kalberer, H. Hengartner, and R. M. Zinkernagel
Formalin inactivation of vesicular stomatitis virus impairs T-cell but not T-help-independent B-cell responses
J. Virol. (1993) 67:3917.

Kündig, T. M., I. Castelmur, M. F. Bachmann, D. Abraham, D. Binder, H. Hengartner, and R. M. Zinkernagel
Fewer protective cytotoxic T cell epitopes than T-helper cell epitopes on vesicular stomatitis virus.
J. Virol. (1993) 67:3680.

Pfeffer, K. , T. Matsuyama, **T. M. Kündig**, A. Wakeham, K. Kishihara, A. Shahinian, K. Wiegmann, P. S. Ohashi, M. Krönke, and T. W. Mak
Mice deficient for the 55 kd tumor necrosis factor receptor are resistant to endotoxic shock, yet succumb to *L. monocytogenes* infection.
Cell (1993) 73:457.

Kishihara, K. , J. Penninger, V. A. Wallace, T. M. Kündig, D. Wakeham, M. Thomas, C. Furlonger, C. J. Paige, and T. W. Mak
Normal B lymphocyte development but impaired T cell maturation in CD45-Exon6 protein

tyrosine phosphatase deficient mice.
Cell (1993) 74:143.

Molina, T. J., M. F. Bachmann, T. M. Kündig, R. M. Zinkernagel, and T. W. Mak
Peripheral T cells in mice lacking p56lck do not express significant anti-viral effector functions.
J. Immunol. (1993) 151:699.

Kündig, T. M., M. F. Bachmann, L. Lefrancois, L. Puddington, H. Hengartner, and R. M. Zinkernagel
Nonimmunogenic tumor cells may efficiently restimulate tumor antigen-specific cytotoxic T cells.
J. Immunol. (1993) 150:4450.

Matsuyama, T. , T. Kimura, M. Kitagawa, K. Pfeffer, T. Kawamaki, N. Watanabe, **T. M. Kündig**, R. Amakawa, K. Kishihara, A. Wakeham, J. Potter, K. Furlonger, A. Narendran, H. Suzuki, P. S. Ohashi, C. J. Paige, T. Taniguchi, and T. W. Mak. Targeted disruption of IRF-1 or IRF-2 results in abnormal type I IFN gene induction and aberrant lymphocyte development.
Cell (1993) 75:83.

Rahemtulla, A. , A. Shahinian, **T. M. Kündig**, R. M. Zinkernagel, and T. W. Mak CD4 negative mice as a model for immunodeficiency.
Phil. Trans. R. Soc. Lond. B (1993) 342:57.

Kündig, T. M., C. P. Kalberer, H. Hengartner, and R. M. Zinkernagel
Vaccination with two different vaccinia recombinant viruses: long term inhibition of secondary vaccination.
Vaccine (1993) 11:1154.

Bachmann, M. F., U. Hoffmann, **T. M. Kündig**, H. Hengartner, and R. M. Zinkernagel
The influence of antigen organization on B cell responsiveness.
Science (1993) 262:1448.

Yeung, R. S. M., J. Penninger, **T. M. Kündig**, Y. Law, K. Yamamoto, N. Kamikawaji, L. Burkly, T. Sasazuki, R. Flavell, P. S. Ohashi, and T. W. Mak
Human CD4-major histocompatibility complex class II (DQw6) transgenic mice in an endogenous CD4/CD8-deficient background: reconstitution of phenotype and human-restricted function
J. Exp. Med. (1994) 180:1911.

Bachmann, M. F., **T. M. Kündig**, G. Freer, Y. Li, C. Y. Kang, D. H. L. Bishop, H. Hengartner, and R. M. Zinkernagel
Induction of protective cytotoxic T cells with non-infectious viral proteins
Eur. J. Immunol. (1994) 24:2228.

Rahemtulla, A. , **T. M. Kündig**, A. Narendran, M. F. Bachmann, M. Julius, C. J. Paige, P. S. Ohashi, R. M. Zinkernagel, and T. W. Mak

Class II restricted T helper cells in CD4 deficient mice.
Eur. J. Immunol. (1994) 24:2213.

Bachmann, M. F., **T. M. Kündig**, C. P. Kalberer, H. Hengartner, and R. M. Zinkernagel
How many B cells are needed to protect against a virus?
J. Immunol. (1994) 152:4235.

Bachmann, M. F., **T. M. Kündig**, H. Hengartner, and R. M. Zinkernagel
Regulation of IgG antibody titers by the amount persisting of immune-complexed antigen.
Eur. J. Immunol. (1994) 24:2567.

Fung-Leung, W. P., **T. M. Kündig**, K. Ngo, J. Panakos, J. De Sousa, E. Wang, P. Ohashi, T. W. Mak, and C. Y. Lau
Reduced thymic maturation but normal effector function of CD8⁺ T cells in CD8 β gene-targeted mice.
J. Exp. Med. (1994) 180:959.

Bachmann, M. F., **T. M. Kündig**, B. Odermatt, H. Hengartner, and R. M. Zinkernagel
Free recirculation of memory B cells versus antigen-dependent differentiation to antibody-forming cells.
J. Immunol. (1994) 153:3386.

Kündig, T. M., M. F. Bachmann, C. DiPaolo, J. L. Simard, M. Battegay, H. Lothar, A. Gessner, K. Kühlke, P. S. Ohashi, H. Hengartner, and R. M. Zinkernagel
Fibroblasts as efficient antigen-presenting cells in lymphoid organs.
Science (1995) 268:1343.

Suzuki, H., **T. M. Kündig**, C. Furlonger, A. Wakeham, E. Timms, T. Matsuyama, R. Schmits, J. L. Simard, P. S. Ohashi, H. Griesser, T. Taniguchi, C. J. Paige, and T. W. Mak
Deregulated T cell activation and autoimmunity in mice lacking interleukin-2 receptor-beta.
Science (1995) 268:1472.

Ally, B. A., T. S. Howley, K. J. McKall-Faienza, **T. M. Kündig**, S. U. Oehen, H. Pircher, R. G. Howley, and P. S. Ohashi
Prevention of autoimmune disease by retroviral-mediated gene therapy.
J. Immunol. (1995) 155:5404.

Kündig, T. M., M. F. Bachmann, S. Oehen, U. W. Hoffmann, J. L. Simard, C. P. Kalberer, H. Pircher, P. S. Ohashi, H. Hengartner, and R. M. Zinkernagel
On the role of antigen in maintaining cytotoxic T-cell memory
Proc. Natl. Acad. Sci. USA (1996) 93:9716.

Yeung, R. S. M., J. M. Penninger, **T. M. Kündig**, W. Khoo, P. S. Ohashi, G. Kroemer, and T. W. Mak
Human CD4 and human MHC class II (DQ6) transgenic mice: a supersensitive model of superantigen induced septic shock.

Eur. J. Immunol. (1996) 26:1074.

Amakawa, R. , A. Hakern, **T. M. Kündig**, T. Matsuyama, J. J. L. Simard, E. Timms, A. Wakeham, H. W. Mittrücker, H. Griesser, H. Takimoto, R. Schmits, A. Shahinian, P. S. Ohashi, J. M. Penninger, and T. W. Mak
Impaired negative selection of T cells in Hodgkin's disease antigen CD30-deficient mice.
Cell (1996) 84:552.

Kündig, T. M., A. Shahinian, K. Kawai, H. W. Mittrücker, E. Sebzda, T. W. Mak, and P. Ohashi
Duration of T cell receptor stimulation determines costimulatory requirements of T cells.
Immunity (1996) 5:41.

Schmits, R. , **T. M. Kündig**, D. M. Baker, G. Schumaker, J. J. L. Simard, G. Duncan, A. Wakeham, A. Shahinian, A. van der Heiden, M. F. Bachmann, P. S. Ohashi, T. W. Mak, and D. D. Hickstein
LFA-1-deficient mice show normal CTL responses to viruses but fail to reject immunogenic tumor.
J. Exp. Med. (1996) 183:1415.

Mittrücker, H. W., A. Shahinian, D. Bouchard, **T. M. Kündig**, and T. W. Mak. Induction of unresponsiveness and impaired T cell expansion and induction of unresponsiveness by staphylococcal enterotoxin B in CD28-deficient mice.
J. Exp. Med. (1996) 183:2481.

Sebzda, E., **T. M. Kündig**, C. T. Thomson, K. Aoki, S. Y. Mak, J. P. Mayer, T. Zamborelli, S. G. Nathanson, and P. S. Ohashi
Mature T cell reactivity altered by peptide agonist that induces positive selection.
J. Exp. Med. (1996) 183:1093.

Bachmann, M.F., E. Sebzda, **T.M. Kündig**, A. Shahinian, D.E. Speiser, T.W. Mak, P.S. Ohashi
T cell responses are governed by avidity and costimulation thresholds.
Eur.J.Immunol. (1996) 26:2017.

Schmits-R; Filmus-J; Gerwin-N; Senaldi-G; Kiefer-F; **Kündig-T**; Wakeham-A; Shahinian-A; Catzavelos-C; Rak-J; Furlonger-C; Zakarian-A; Simard-JJ; Ohashi-PS; Paige-CJ; Gutierrez-Ramos-JC; Mak-TW
CD44 regulates hematopoietic progenitor distribution, granuloma formation, and tumorigenicity.
Blood. 1997 Sep 15; 90(6): 2217-33

Dommann-SN; Dommann-Scherrer-CC; Ziegler-T; Meyer-J; Trueb-RM; **Kündig-T**; Panizzon-R; Burg-G
Expression of intercellular adhesion molecule 3 (CDw50) on endothelial cells in cutaneous lymphomas. A comparative study between nodal and cutaneous lymphomas.
Am-J-Dermatopathol. 1997 Aug; 19(4): 391-5

Kündig-TM; Trueb-RM; Krasovec-M.
Lupus profundus/panniculitis
Dermatology. 1997; 195(1): 99-101

Zinkernagel-RM; Ehl-S; Aichele-P; Oehen-S; **Kündig-T**; Hengartner-H
Antigen localisation regulates immune responses in a dose- and time-dependent fashion: a geographical view of immune reactivity.
Immunol-Rev. 1997 Apr; 156: 199-209

Krause-MH; Torricelli-R; **Kündig-T**; Trueb-RM; Hafner-J
[Dapsone in granulomatous rosacea]
Hautarzt. 1997 Apr; 48(4): 246-8

Wallace-VA; Penninger-JM; Kishihara-K; Timms-E; Shahinian-A; Pircher-H; **Kündig-TM**;
Ohashi-PS; Mak-TW
Alterations in the level of CD45 surface expression affect the outcome of thymic selection.
J-Immunol. 1997 Apr 1; 158(7): 3205-14

Kozieradzki-I; **Kündig-T**; Kishihara-K; Ong-CJ; Chiu-D; Wallace-VA; Kawai-K; Timms-E;
Ionescu-J; Ohashi-P; Marth-JD; Mak-TW; Penninger-JM
T cell development in mice expressing splice variants of the protein tyrosine phosphatase CD45.
J-Immunol. 1997 Apr 1; 158(7): 3130-9

Bachmann-MF; **Kündig-TM**; Hengartner-H; Zinkernagel-RM
Protection against immunopathological consequences of a viral infection by activated but not resting cytotoxic T cells: T cell memory without "memory T cells"?
Proc-Natl-Acad-Sci-U-S-A. 1997 Jan 21; 94(2): 640-5

Bachmann-MF; **Kündig-TM**; Speiser-DE; McKall-Faienza-K; Kishara-K; Mak-TW; Ohashi-PS
T-cell-independent antiviral B cell responses in CD45-deficient mice.
Cell-Immunol. 1997 Jan 10; 175(1): 12-5

Mittrucker-HW; Matsuyama-T; Grossman-A; **Kündig-TM**; Potter-J; Shahinian-A; Wakeham-A;
Patterson-B; Ohashi-PS; Mak-TW
Requirement for the transcription factor LSIRF/IRF4 for mature B and T lymphocyte function.
Science. 1997 Jan 24; 275(5299): 540-3

Schmits-R; Filmus-J; Gerwin-N; Senaldi-G; Kiefer-F; **Kündig-T**; Wakeham-A; Shahinian-A;
Catzavelos-C; Rak-J; Furlonger-C; Zakarian-A; Simard-JJ; Ohashi-PS; Paige-CJ; Gutierrez-Ramos-JC; Mak-TW
CD44 regulates hematopoietic progenitor distribution, granuloma formation, and tumorigenicity.
Blood. 1997 Sep 15; 90(6): 2217-33

Dommann-SN; Dommann-Scherrer-CC; Ziegler-T; Meyer-J; Trueb-RM; **Kündig-T**; Panizzon-R; Burg-G

Expression of intercellular adhesion molecule 3 (CDw50) on endothelial cells in cutaneous lymphomas. A comparative study between nodal and cutaneous lymphomas.
Am-J-Dermatopathol. 1997 Aug; 19(4): 391-5

Kündig-TM; Trueb-RM; Krasovec-M.
Lupus profundus/panniculitis
Dermatology. 1997; 195(1): 99-101

Krause-MH; Torricelli-R; **Kündig-T**; Trueb-RM; Hafner-J
[Dapsone in granulomatous rosacea]
Hautarzt. 1997 Apr; 48(4): 246-8

McKall-Faienza-KJ; Kawai-K; **Kündig-TM**; Odermatt-B; Bachmann-MF; Zakarian-A; Mak-TW; Ohashi-PS
Absence of TNFRp55 influences virus-induced autoimmunity despite efficient lymphocytic infiltration.
Int-Immunol. 1998; 10(4): 405-12

Maloy-K, Eerdmann-I, Basch-V, Sierro-S, Kramps-T, Zinkernagel-RM, Oehen-S, **Kündig-TM**
Intralymphatic immunization enhances DNA vaccination
Proc. Natl. Acad. Sci. USA 2001; 98: 3299-3203

Decreased intraindividual HLA class I expression is due to reduced transcription in advanced melanoma and does not correlate with HLA-G expression
Willers, J., Urosevic, M., Laine, E., Geertsen, R., **Kündig, T.M.** Burg, G., Dummer, R.
J Invest Dermatol 117:000-000, 2001

Urtikaria und Quincke-Oedem

T.M. Kündig

Allergolist 2001

Therapie der chronischen Urtikaria und der rezidivierenden Quincke-Oedeme

T.M. Kündig

Therapeutische Umschau, Band 58, Heft 5, 2001, Seite 321-324

Verfahren zur Steigerung der Immunogenität von Impfstoffen

T.M. Kündig

Praxis, Schweizerische Rundschau für Medizin, 89 (37), 1477-84

Fucosyltransferase-VII deficient mice with defective E-, P- and L-selectin ligands show impaired CD4⁺ and CD8⁺ T cell migration into the skin but normal extravasation into visceral organs
Erdmann, I., Scheidegger, E.P., Koch, F., Heinzerling, L., Odermatt, B., Burg, G., Lowe, J.B.,
and **Kündig, T.M.**

J Immunol. 2002 Mar 1;168(5):2139-46.

Critical role of activation of antigen presenting cells in priming of cytotoxic T cell responses after vaccination with virus-like particles

Tazio Storni, Franziska Lechner, Iris Erdmann, Thomas Bächli, Andrea Jegerlehner, Tilman Dumrese, **Thomas M. Kündig**, Christiane Ruedl, Martin F. Bachmann

J. Immunol. 2002, 15; 168 (&):2880-6

The use of anti-T-cell receptor-Vbeta antibodies for the estimation of treatment success and phenotypic characterization of clonal T-cell populations in cutaneous T-cell lymphomas.

Schwab C, Willers J, Niederer E, Lucwig E, **Kündig T**, Grob P, Burg G, Dummer R

Br J Haematol. 2002 Sep;118(4):1019-26.

A molecular assembly system that renders antigens of choice highly repetitive for induction of protective B cell responses.

Jegerlehner A, Tissot A, Lechner F, Sebbel P, Erdmann I, **Kündig T**, Bächli T, Storni T, Jennings G, Pumpens P, Renner WA, Bachmann MF

Vaccine. 2002 Aug 19;20(25-26):3104-12.

Proliferation of CD30+ T-Helper 2 Lymphoma Cells Can Be Inhibited by CD30 Receptor Cross-Linking with Recombinant CD30 Ligand.

Willers J, Dummer R, Kempf W, **Kündig T**, Burg G, Kadin ME

Clin Cancer Res. 2003 Jul;9(7):2744-54.

Publications (Reviews):

Bachmann, M. F. and T. M. Kündig

In vitro vs. in vivo assays for the assessment of T- and B-cell function.

Curr. Opin. Immunol. (1994) 6:320.

Zinkernagel, R. M., M. F. Bachmann, T. M. Kündig, S. Oehen, and H. Hengartner

On immunological memory.

Annu. Rev. Immunol. (1996) 14:333.

Kündig, T. M., M. F. Bachmann, P. S. Ohashi, H. Pircher, H. Hengartner, and R. M.

Zinkernagel

On T cell memory: arguments for antigen dependence.

Immunol. Rev. (1996) 150:63.

Kündig, T.M.

30 Jahre IgE.

Labolife (1997)

Zinkernagel-RM; Ehl-S; Aichele-P; Oehen-S; Kündig-T; Hengartner-H

Antigen localisation regulates immune responses in a dose- and time-dependent fashion: a geographical view of immune reactivity.

Immunol-Rev. 1997 Apr; 156: 199-209

Kündig, T.M.

Immunity and cancer

Advances in Biology and Medicine

Plenum publishers, in press, New York, USA

EXHIBIT 2

Study ZU-GTP-001




EXHIBIT 3

Nasal Provocation Symptom Scores






EXHIBIT 4


Subcutaneous



Intralymphatic



Subcutaneous



Intralymphatic




EXHIBIT 5

IgG4 against Grass (Ph. Pratensae)

